

## CASE REPORT

## A case of maxillary sarcoma in a chimpanzee (*Pan troglodytes*)

M. Fujisawa<sup>1,2</sup>, T. Usono<sup>1</sup>, E. Nogami<sup>1</sup>, M. Hirosawa<sup>1</sup>, N. Morimura<sup>1</sup>, A. Saito<sup>1</sup>, M. Seres<sup>3</sup>, M. Teramoto<sup>1</sup>, K. Nagano<sup>1</sup>, Y. Mori<sup>1</sup>, H. Uesaka<sup>1</sup>, K. Nasu<sup>1</sup>, M. Tomonaga<sup>3</sup>, G. Idani<sup>1</sup>, S. Hirata<sup>3</sup>, T. Tsuruyama<sup>4</sup> & K. Matsubayashi<sup>2</sup>

1 Kumamoto Sanctuary, Wildlife Research Center, Kyoto University, Uki, Japan

2 Center for Southeast Asian Studies, Kyoto University, Kyoto, Japan

3 Primate Research Institute, Kyoto University, Inuyama, Japan

4 Department of Diagnostic Pathology, Kyoto University Hospital, Kyoto, Japan

### Keywords

aging – ape – care – malignant neoplasm

### Correspondence

Michiko Fujisawa, Center for Southeast Asian Studies, Kyoto University, 46, Shimoadachi-cho, Yosida, Sakyo-ku, Kyoto-shi, Kyoto 606-8501, Japan.

Tel.: 075 753 7302;

fax: 075 753 7350;

e-mail: mfujisaw@cseas.kyoto-u.ac.jp

Accepted October 14, 2013.

### Abstract

Oral malignancy is rare in chimpanzees. A 34-year-old female chimpanzee (*Pan troglodytes*) at Kumamoto Sanctuary, Japan, had developed it. Treatment is technically difficult for chimpanzees while malignant neoplasm is seemingly rising in captive populations. Widespread expert discussion, guidelines for treatment, especially for great apes in terminal stages is urgently needed.

### Case report

Genetically chimpanzees are the closest living relatives of humans. We were diverged almost 7 million years ago, and the changes in disease conditions could have occurred in conjunction with evolutionary changes. As our life expectancies increased, the later-onset diseases such as malignant neoplasm, arteriosclerotic diseases, and dementia have also increased, thus becoming a serious issue. These diseases are dependent on the lifestyle and aging; they could be referred to as age-dependent diseases.

In contrast, very little is known about the nature of age-dependent diseases in chimpanzees. The question remains whether the age-dependent diseases are exclusively human or simply undiagnosed in chimpanzees. For example, malignancy is extremely rare in chimpanzees [4]. Previous studies have shown only few examples of such diseases in chimpanzees [2, 5].


This is the first report of oral sarcoma in a chimpanzee. The case occurred in an estimated 34-year-old wild-born female. She was utilized in hepatitis C virus (HCV) infection research at another facility between 1979 and

1987. Persistent HCV infection was observed without other specific notations.


The subject's right cheek had started swelling in February 2011. We suspected bacterial infection of the dental root. Antibiotics were administered; however, no improvement was observed (Fig. 1). Blood analysis showed slight increases in white blood cell, C-reactive protein, and  $\gamma$ -glutamyl transpeptidase levels and decrease in albumin level. No other noteworthy observations were made. Her weight was 43 kg.

Figure 2 shows histopathological image of the lesion. Abnormal undifferentiated cells including spindle cells and adipose-like cells proliferated papillary or focally. Increased N/C ratio and multiple images of mitosis showed this tumor was highly malignant. By immunohistological stain, vimentin was positive, cytokeratin was partially slightly positive. On X-ray examination maxilla invasion was identified; however, no pulmonary metastasis could be observed. Cervical lymph nodes were not swollen.

The common treatment for sarcoma in humans is mainly surgical resection with radiation or chemotherapy.



**Fig. 1** (A) Swelling around the nose. (B) Tumor inside the mouth



**Fig. 2** (A, B) Proliferation of spindle cells and adipose-like cells. (C, D) Proliferation of spindle cells with mitosis. (E, F) Sarcoma like change with atypical cells.

In this case, because extensive resection was required, the reconstruction of the oral cavity must be considered. Surgical resection and radiation could not be options due to technical difficulties. Chemotherapy was avoided as the side effects were thought to be too severe in proportion to the potential results. It was also difficult to give her injection due to risking anesthesia on a daily basis. A conservative palliative treatment was chosen instead during the remaining progression of the disease.

The tumor reached the end of hard palate, began to construct the pharynx. However, food intake was still achieved by altering the texture of food. A timetable was created incorporating contact time with other chimpanzees, with staff, and time spent alone. Gradually, the time spent being recumbent increased; however, she would still rise to interact with staff. Breathing difficulty appeared except when lying on her right side. On August 15, she had eventually attempted to roll over

and reached with her arms and legs out to staff. She died at 9:30 am on August 17. Euthanasia was debated during the course, but was not performed because she was still able to ingest food and did not appear to be in great distress.

When she died her weight was 26 kg. Pathological autopsy showed that the tumor had already invaded her maxilla. The mass had protruded into the oral cavity, and it was largely necrotic and ulcerated, but the tumor had not yet progressed into the orbital cavity. Despite the tumor extended almost to cover the entire palate, it did not reach the pharynx. The mandible and cervical lymph nodes were largely swollen, and the right cervical lymph nodes entwined the carotid artery. Multiple metastatic lesions were identified in lymph nodes of the pulmonary hilum and of posterior mediastinum, in lungs, and in diaphragm. The right lung showed poor aeration with the lower lung atelectasis. No pleural effusion was observed. No intraperitoneal spread was detected.

Pathological diagnosis: (i) maxillary sarcoma with multiple metastases, (ii) respiratory failure due to metastases.

There are very few reports of malignant tumors in chimpanzees. Till recently, a number of hypotheses have been raised, including claims that malignant tumors are simply remain undiagnosed [5] or that because apoptosis suppression difference between human and chimpanzees, there is a known relationship between suppression of apoptosis and increased risk of the onset of malignancy [3, 11]. Recent studies also have shown multiple genetic differences associated with malignancies [1, 7, 13, 14]. From these findings, it is now understood that currently, there are very few identified genetic predispositions for malignant neoplasm in chimpanzees.

The ratio of chimpanzees that actually reaches an advanced age is low [8]. Although the frequency of occurrence of malignancy because of gene restoration anomalies increases with aging, the frequency of malignancy remains low in chimpanzees due to the shorter life

span. This can be a possible explanation for the minimal number of age-related diseases that are observed in chimpanzees.

The case subject had persistent HCV infection. In humans, oral squamous-cell carcinoma is considered as a complication of HCV infection [6, 9, 10, 12]. Chimpanzees with a history of being utilized in hepatitis research are numerous in Japan and in several countries, including USA. Although cases of hepatic carcinoma have been reported, thus far, there have been no reports of oral malignancy.

As outlined above, the possibility is that HCV infection is partially responsible for the oral sarcoma. As the subject was not young, the potential for gene restoration mistakes to occur at a higher rate along with advancement in age must also be taken into consideration.

Noteworthy, that the subject was able to ingest food orally till the end; further, pain did not seem to appear. The subject was able to live out the remainder of her life in relative comfort during the terminal phase of the disease due to devoted staff until her very last day. The actual treatment of a chimpanzee during the terminal phase of disease may be decided on a case-to-case basis. The number of aging chimpanzees in captivity is increasing, and so we have to consider the need for terminal care cases will also increase. Therefore, we think the necessity for a widespread discussion, regarding these issues is inevitable among other chimpanzee holding facilities.

### Acknowledgments

We thank Drs Yasunori Sumi and Zenzo Isogai of National Center for Geriatrics and Gerontology Hospital for providing valuable suggestion on treating malignant tumor in oral cavity and skin. This study was financially supported by the following grants: MEXT 20002001, 20040001, 22659141, and WISH Grant and Human Evolution Grant to the Primate Research Institute, Kyoto University.

### References

- Ahmed J, Günther S, Möller F, Preissner R: A structural genomics approach to the regulation of apoptosis: chimp vs. human. *Genome Inform* 2007; **18**:22–34.
- Amyx HL, Salazar AM, Newsome DA, Gibbs CJ Jr, Gajdusek DC: Nasopharyngeal carcinoma with intracranial extension in a chimpanzee. *J Am Vet Med Assoc* 1982; **181**:1425–6.
- Arora G, Polavarapu N, McDonald JF: Did natural selection for increased cognitive ability in humans lead to an elevated risk of cancer? *Med Hypotheses* 2009; **73**:453–6.
- Bailey J: An examination of chimpanzee use in human cancer research. *Alter Lab Anim* 2009; **37**:399–416.
- Brown SL, Anderson DC, Dick Jr EJ, Rodolfo G-M, AnaPatricia G, Gene BH: Neoplasia in the Chimpanzee (*Pan spp.*). *J Med Primatol* 2009; **38**:137–44.
- Carrozzo M: Oral diseases associated with hepatitis C virus infection.

- Part 2: lichen planus and other diseases. *Oral Dis* 2008; **14**:217–28.
- 7 Calarco JA, Xing Y, Cáceres M, Clarco JP, Xiao X, Pan Q, Lee C, Preuss TM, Blencowe BJ: Global analysis of alternative splicing differences between humans and chimpanzees. *Genes Dev* 2007; **21**:2963–75.
- 8 Finch CE: Evolution of the human lifespan and disease of aging: role of infection, inflammation, and nutrition. *Proc Natl Acad Sci* 2010; **107**:1718–24.
- 9 Gandolfo S, Richiardi L, Carrozzo M, Broccoletti R, Carbone M, Pagano M, Vestita C, Rosso S, Merletti F: Risk of oral squamous cell carcinoma in 402 patients with oral lichen planus: a follow-up study in an Italian population. *Oral Oncol* 2004; **40**:77–83.
- 10 Grossmann SM, Teixeira R, de Aguiar MC, de Moura MD, do Carmo MA: Oral mucosal conditions in chronic hepatitis C Brazilian patients: a cross-sectional study. *J Public Health Dent* 2009; **69**:168–75.
- 11 Lowe SW, Lin AW: Apoptosis in cancer. *Carcinogenesis* 2000; **21**:485–95.
- 12 Nagao Y, Sata M, Noguchi S, Seno'o T, Kinoshita M, Kameyama T, Ueno T: Detection of hepatitis C virus RNA in oral lichen planus and oral cancer tissues. *J Oral Pathol Med* 2000; **29**:259–66.
- 13 Wallace DC, Brown MD, Lott MT: Mitochondrial DNA variation in human evolution and disease. *Gene* 1999; **238**:211–30.
- 14 Wallace DC: A mitochondrial paradigm of metabolic and degenerative diseases, aging, and cancer: a dawn for evolutionary medicine. *Annual Rev Genet* 2005; **39**:359–407.